


Johnson & Johnson Consumer Inc.

-PE Product Labels –



Drug Facts	
Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr	Cough suppressant
Phenylephrine HCl	Nasal decongestant
Guafenesin	Expectorant

Drug Facts (continued)	
temporarily reduces fever	
Warnings	
Liver warning: This product contains acetaminophen. Severe liver damage	

Drug Facts (continued)	
If a skin reaction occurs, stop use and seek medical help right away.	
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever,	

Dextromethorphan HBr 20 mg.....Cough suppressant
Guaifenesin 400 mg.....Expectorant
Phenylephrine HCl 10 mg.....Nasal decongestant

Uses

- temporarily relieves the following cold/flu symptoms:
 - minor aches and pains
 - headache
 - nasal congestion
 - sore throat
 - cough
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression,

Drug Facts (continued)

psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

Drug Facts (continued)

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dose

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days

Drug Facts (continued)

- redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- do not take more than directed (see overdose warning)
- mL = milliliter
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

adults and children 12 years and over

- take 30 mL in the dosing cup provided every 4 hours while symptoms last
- do not take more than 150 mL in 24 hours, unless directed by a doctor

children under 12 years

ask a doctor

Drug Facts (continued)

Other information

- each 30 mL contains: sodium 10 mg
- store between 20-25°C (68-77°F).
- do not use if neck band imprinted with "WARNING SENSATION" or foil inner seal imprinted with "TYLENOL" is broken or missing

Inactive ingredients

anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, FD&C yellow no. 6, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

Drug Facts (continued)

Questions or comments?

call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

OPEN TO READ DRUG FACTS
(Warnings, Directions...)

PEEL




Do not use if neck band imprinted with "TYLENOL COLD MAX" or foil inner seal imprinted with "SAFETY SEAL®" is broken or missing.

TYLENOL®
COLD MAX

How can we help?
1-877-895-3665

SNAP THIS CODE
OR TEXT TYCOLD7
TO 87715

Acetaminophen
(325 mg per 15 mL)
Dextromethorphan HBr



MSG & DATA RATES MAY APPLY
TEXT HELP FOR HELP
TEXT STOP FOR STOP



(10 mg per 15 mL)
Doxylamine succinate
(6.25 mg per 15 mL)
Phenylephrine HCl
(5 mg per 15 mL)

8 fl oz (240 mL)

Made in Canada

Distributed by:

JOHNSON & JOHNSON CONSUMER INC.

McNeil Consumer Healthcare Division

Fort Washington, PA 19034 USA

© J&JCI 2018

30042166

Drug Facts

Active ingredients (in each 15 mL) Purpose

Acetaminophen 325 mg Pain reliever/
fever reducer

Dextromethorphan HBr 10 mg Cough suppressant
Doxylamine succinate 6.25 mg Antihistamine
Phenylephrine HCl 5 mg Nasal decongestant

Uses

■ temporarily relieves these common cold/flu symptoms:

- minor aches and pains
- headache
- sore throat
- nasal congestion
- runny nose and sneezing
- cough
- sinus congestion and pressure

Drug Facts (continued)

- helps clear nasal passages
- relieves cough to help you sleep
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions.

Symptoms may include:
■ skin reddening

Drug Facts (continued)

- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription

Drug Facts (continued)

monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged

Drug Facts (continued)

prostate gland

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not exceed recommended dose
- excitability may occur,

Drug Facts (continued)

especially in children

- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

Drug Facts (continued)

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ **do not take more than directed** (see overdose warning)

■ mL = milliliter

■ use only enclosed dosing cup ►

Drug Facts (continued)

designed for use with this product. Do not use any other dosing device.

adults
and children
12 years
and over

■ take 30 mL in the dosing cup provided every 4 hours while symptoms last

■ do not take more than 150 mL in 24 hours, unless directed by a doctor

children under
12 years

ask a doctor

Other information

■ each 15 mL contains: sodium 5 mg

■ store between 20-25°C (68-77°F). Do not refrigerate.

■ **do not use if neck band imprinted with "TYLENOL COLD MAX" or** ►

Drug Facts (continued)

foil inner seal imprinted with "SAFETY SEAL®" is broken or missing

Inactive ingredients

alcohol, anhydrous citric acid, FD&C blue no. 1, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

Questions or comments?

call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org



OPEN TO READ DRUG FACTS
(Warnings, Directions...)

PEEL

Do not use if neck band imprinted with
"TYLENOL COLD+FLU+ COUGH" or foil
inner seal imprinted with "SAFETY
SEAL®" is broken or missing.

DO NOT USE WITH OTHER MEDICINES
CONTAINING ACETAMINOPHEN



Drug Facts

Active ingredients Purpose

Acetaminophen 325 mg Pain reliever/
fever reducer

Dextromethorphan HBr 10 mg.....Cough suppressant
Doxylamine Succinate 6.25 mg.....Antihistamine
Phenylephrine HCl 5 mg.....Nasal decongestant

Uses

- temporarily relieves these common cold/flu symptoms:
 - minor aches and pains
 - headache
 - nasal congestion
 - runny nose and sneezing
 - cough
 - sinus congestion and pressure
 - helps clear nasal passages
- sore throat

Drug Facts (continued)

- relieves cough to help you sleep
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional)

Drug Facts (continued)

conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as

Drug Facts (continued)

- emphysema or chronic bronchitis
- glaucoma
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not exceed recommended dose
- excitability may occur, especially in children
- marked drowsiness may occur

Drug Facts (continued)

- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
 - pain, nasal congestion or cough gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts
- These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Drug Facts (continued)

adults take 30 mL in the dosing cup provided every 4

Drug Facts (continued)

is broken or missing

TYLENOL®
COLD + FLU + COUGH
NIGHT

Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
- mL = milliliter
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

and children 12 years and over	cup provided every 4 hours while symptoms last
children under 12 years	ask a doctor

Other information

- each 15 mL contains: sodium 5 mg
- store between 20-25°C (68-77°F). Do not refrigerate.
- do not use if neck band imprinted with "TYLENOL COLD+FLU+COUGH" or foil inner seal imprinted with "SAFETY SEAL®"

Inactive ingredients

acetic acid, alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

Questions or comments?

call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

How can we help?

1-877-895-3665

SNAP THIS CODE OR TEXT TYCOLD7 TO 87715

MSG & DATA RATES MAY APPLY
 TEXT HELP FOR HELP
 TEXT STOP FOR STOP



Acetaminophen (325 mg per 15 mL)
 Dextromethorphan HBr (10 mg per 15 mL)
 Doxylamine succinate (6.25 mg per 15 mL)
 Phenylephrine HCl (5 mg per 15 mL)
 8 fl oz (240 mL)


Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
 McNeil Consumer Healthcare Division
 Fort Washington, PA 19034 USA
 © J&JCI 2017

30040372

NDC 50580-403-08 30049843

Warming Honey Lemon Flavor

TYLENOL®
 FOR ADULTS



Drug Facts

Active ingredients Purpose (in each 30 mL)

Acetaminophen 650 mg.....	Pain reliever/ fever reducer
Dextromethorphan HBr 20 mg.....	Cough suppressant
Guaifenesin 400 mg.....	Expectorant
Phenylephrine HCl 10 mg.....	Nasal decongestant

Uses

- temporarily relieves the following cold/flu

Drug Facts (continued)

- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day

symptoms:

- minor aches and pains
- headache
- nasal congestion
- sore throat
- cough
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

- 3 or 12 alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression,

Drug Facts (continued)

psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

Drug Facts (continued)

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dose

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days

Drug Facts (continued)

- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)**Drug Facts (continued)**

Directions

- do not take more than directed (see overdose warning)
- mL = milliliter
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

adults
and
children
12 years
and over

- take 30 mL in the dosing cup provided every 4 hours while symptoms last
- do not take more than 150 mL in 24 hours, unless directed by a doctor

children
under
12 years

ask a doctor

Other information

- each 30 mL contains: sodium 10 mg
- store between 20-25°C (68-77°F).
- do not use if neck band imprinted with "WARMING SENSATION" or foil inner seal imprinted with "TYLENOL" is broken or missing

Inactive ingredients

anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, FD&C yellow no. 6, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

Drug Facts (continued)**Questions or comments?**

call 1-877-895-3665 (toll-free) or
215-273-8755 (collect)

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

TYLENOL®
COLD + FLU SEVERE

How can we help?

1-877-895-3665

Acetaminophen
(650 mg per 30 mL)
Dextromethorphan HBr
(20 mg per 30 mL)
Guaifenesin
(400 mg per 30 mL)
Phenylephrine HCl
(10 mg per 30 mL)

8 fl oz (240 mL)

Distributed by:

JOHNSON & JOHNSON CONSUMER INC.

McNeil Consumer Healthcare Division

Fort Washington, PA 19034 USA

© J&JCI 2021

30049845



OPEN TO READ DRUG FACTS
(Warnings, Directions...)

PEEL

Do not use if neck band imprinted with
"TYLENOL COLD+FLU+ COUGH" or foil
inner seal imprinted with "SAFETY
SEAL®" is broken or missing.

DO NOT USE WITH OTHER MEDICINES
CONTAINING ACETAMINOPHEN



Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg Pain reliever/
fever reducer
Dextromethorphan HBr 10 mg... Cough suppressant
Doxylamine Succinate 6.25 mg.....Antihistamine
Phenylephrine HCl 5 mg.....Nasal decongestant

Uses

- temporarily relieves these common cold/flu symptoms:
 - minor aches and pains
 - headache
 - nasal congestion
 - runny nose and sneezing
 - cough
 - sinus congestion and pressure
 - helps clear nasal passages
 - sore throat

Drug Facts (continued)

- relieves cough to help you sleep
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional

Drug Facts (continued)

conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as

Drug Facts (continued)

- emphysema or chronic bronchitis
- glaucoma
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not exceed recommended dose
- excitability may occur, especially in children
- marked drowsiness may occur

Drug Facts (continued)

- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
 - pain, nasal congestion or cough gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts
- These could be signs of a serious condition.

Drug Facts (continued)

Drug Facts (continued)

Drug Facts (continued)

TYLENOL

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
- mL = milliliter
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

adults
and
children
12 years
and over

- take 30 mL in the dosing cup provided every 4 hours while symptoms last
- do not take more than 150 mL in 24 hours, unless directed by a doctor

children
under
12 years

ask a doctor

Other information

- each 15 mL contains: sodium 5 mg
- store between 20-25°C (68-77°F). Do not refrigerate.
- do not use if neck band imprinted with "TYLENOL COLD+FLU+COUGH" or foil inner seal imprinted with "SAFETY SEAL®"

is broken or missing

Inactive ingredients

acetic acid, alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

Questions or comments?

call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

COLD + FLU + COUGH NIGHT

How can we help?

1-877-895-3665

SNAP THIS CODE
OR TEXT TYCOLD7
TO 87715

MSG & DATA RATES MAY APPLY
TEXT HELP FOR HELP.
TEXT STOP FOR STOP



Acetaminophen
(325 mg per 15 mL)
Dextromethorphan HBr
(10 mg per 15 mL)
Doxylamine succinate
(6.25 mg per 15 mL)
Phenylephrine HCl
(5 mg per 15 mL)

8 fl oz (240 mL)

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
 McNeil Consumer Healthcare Division
 Fort Washington, PA 19034 USA
 © J&JCI 2017

30040372

NDC 50580-403-08 30049843

Warming Honey Lemon Flavor

TYLENOL®
FOR ADULTS



Drug Facts

Active ingredients Purpose (in each 30 mL)

Acetaminophen 650 mg.....	Pain reliever/ fever reducer
Dextromethorphan HBr 20 mg.....	Cough suppressant
Guaifenesin 400 mg.....	Expectorant
Phenylephrine HCl 10 mg.....	Nasal decongestant

Drug Facts (continued)

- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing

Uses

- temporarily relieves the following cold/flu symptoms:
 - minor aches and pains
 - headache
 - nasal congestion
 - sore throat
 - cough
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

acetaminophen

- 3 or more alcoholic drinks every day while using this product
- Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
 - blisters
 - rash

Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression,

Drug Facts (continued)

psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

Drug Facts (continued)

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dose

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days

Drug Facts (continued)

- redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)**Directions**

- do not take more than directed (see overdose warning)
- mL = milliliter
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

adults
and
children
12 years
and over

- take 30 mL in the dosing cup provided every 4 hours while symptoms last
- do not take more than 150 mL in 24 hours, unless directed by a doctor

children
under
12 years

ask a doctor

Drug Facts (continued)**Other information**

- each 30 mL contains: sodium 10 mg
- store between 20-25°C (68-77°F).
- do not use if neck band imprinted with "WARMING SENSATION" or foil inner seal imprinted with "TYLENOL" is broken or missing

Inactive ingredients

anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, FD&C yellow no. 6, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

Drug Facts (continued)**Questions or comments?**

call 1-877-895-3665 (toll-free) or
215-273-8755 (collect)

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

TYLENOL®
COLD + FLU SEVERE

How can we help?

☎ 1-877-895-3665

Acetaminophen
(650 mg per 30 mL)
Dextromethorphan HBr
(20 mg per 30 mL)
Guaifenesin
(400 mg per 30 mL)
Phenylephrine HCl
(10 mg per 30 mL)

8 fl oz (240 mL)

Distributed by:

JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
© J&JCI 2021

30049845

Diphenhydramine HCl 25mg | Antihistamine
Phenylephrine HCl 10mg | Nasal Decongestant

ALLERGY PLUS

The Benadryl logo, featuring the word "Benadryl" in white, bold, sans-serif font, set against a blue background with a pink border.

NDC 50580-415-01

EFFECTIVE ALLERGY RELIEF WHEN YOU NEED IT!®

ALLERGY PLUS



Purposes

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies;

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- nasal congestion

- temporarily relieves these symptoms due to the common cold:
- runny nose ■ sneezing ■ nasal congestion

- temporarily relieves sinus congestion and pressure

Do not use

- to make a child sleepy
- with any other product containing dphenhydramine, even one used on skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- do not exceed recommended dose
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

adults and children
12 years and over

- take 1 tablet every 4 hours
- do not take more than 6 tablets in 24 hours

[ask a doctor](#)

- store between 20-25°C (68-77°F).
- do not use if carton is opened or if blister unit is broken

Inactive ingredients carnauba wax, FD&C blue no. 1 aluminum lake, magnesium stearate, microcrystalline cellulose, modified starch, polyethylene glycol, polyvinyl alcohol, powdered cellulose, pregelatinized starch, sodium starch glycolate, talc, titanium dioxide

call 1-888-217-2117 (toll-free) or 215-273-8755 (collect)



3 0045-0557-26

30037630/
116786


ALLERGY PLUS CONGESTION

Made in Italy
Distributed by: J&JCI
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
@J&JCI 2016

EFFECTIVE ALLERGY RELIEF WHEN YOU NEED IT![®]

ALLERGY PLUS







ALLERGY *PLUS* CONGESTION

How can we help?
1-877-717-2824
benadryl.com

NDC 50580-341-01



ALLERGY *PLUS* CONGESTION

Diphenhydramine HCl / antihistamine
Phenylephrine HCl / nasal decongestant
oral solution

 **4 Hours/Dose**

RELIEF OF:

- ✓ Runny Nose
- ✓ Sneezing
- ✓ Itchy, Watery Eyes
- ✓ Itchy Throat or Nose
- ✓ Sinus Congestion
- ✓ Stuffy Nose

Alcohol Free
Sugar Free



Actual Size



Tips for Outdoor Allergens:

- Avoid being outside when the pollen count is high, typically in the morning
- Shower and change clothing after spending time outdoors



Tips for Indoor Allergens:

- Wash sheets and blankets in hot water and use the dryer weekly





Drug Facts (continued)

- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help.

CONGESTION**Drug Facts**

Active ingredients (in each 5 mL)	Purposes
Diphenhydramine HCl 12.5 mg.....	Antihistamine
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
 - nasal congestion
 - stuffy nose
- temporarily relieves these symptoms due to the common cold:
 - runny nose
 - sneezing
 - nasal congestion
 - stuffy nose
- temporarily relieves sinus congestion and pressure

Warnings**Do not use**

- to make a child sleepy
- with any other product containing diphenhydramine, even one used on skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers**When using this product**

- do not exceed recommended dose
- marked drowsiness may occur
- avoid alcoholic drinks

Directions

- find right dose on chart below
- mL = milliliter
- take every 4 hours
- do not take more than 6 doses in 24 hours

Age (yr)	Dose (mL)
children under 4 years	do not use
children 4 to 5 years	do not use unless directed by a doctor
children 6 to 11 years	5 mL
adults and children 12 years and over	10 mL

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL contains: sodium 10 mg
- store between 20-25°C (68-77°F)
- do not use if carton tape or bottle wrap imprinted with "Sealed For Your Safety" is broken or missing

Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments?

call 1-877-717-2824 (toll-free) or 215-273-8755 (collect)

Distributed by:

JOHNSON & JOHNSON CONSUMER INC.

McNeil Consumer Healthcare Division

Fort Washington, PA 19034 USA

©J&JCI 2019

30046178

For more information, visit:

www.benadryl.com

Drug Facts (continued)

Directions

- find right dose on chart below
- mL = milliliters
- repeat dose every 4 hours
- do not give more than 6 times in 24 hours

Age (yr)	Dose (mL)
under 4 years	do not use
4 to 5 years	5 mL
6 to 11 years	10 mL

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other Information

- each 5 mL contains: sodium 14 mg
- store between 20-25°C (68-77°F). Protect from light.
- Store in outer carton until contents are used.
- do not use if carton tape or bottle wrap imprinted with "Sealed For Your Safety" is broken or missing

Inactive Ingredients

anhydrous citric acid, carbocymethylcellulose sodium, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments?

call 1-888-217-2117 (toll-free) or 215-273-8755 (collect)

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

3 0045-0538-04 8

Actual Size

available behind the counter

Other Products Formulated To Relieve Your Child's Symptoms

Tips for Relieving Stuffy nose and Cough symptoms

Make sure they drink lots of fluids, especially warm drinks

Show them how to breathe in moist air from a hot shower, humidifier or vaporizer

Give them a non-medicated cough drop to help soothe their cough

When treating your child's symptoms, be sure to use the dosing cup to properly dose medication

children's
SUDAFED®
PE

How can we help?

1-888-217-2117
sudafed.com

children's
SUDAFED®
PE
non-drowsy

COLD + COUGH

Phenylephrine HCl • Dextromethorphan HBr
Nasal Decongestant • Cough Suppressant
Oral Solution

RELIEF OF
✓ Stuffy Nose
✓ Sinus Pressure
✓ Cough

Grape
FLAVOR LIQUID
ALCOHOL &
SUGAR FREE

4 fl oz (118 mL)

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19084 USA
© J&JCI 2010

30047105

children's
SUDAFED®
PE
non-drowsy
COLD + COUGH

Drug Facts

Active Ingredients

(In each 5 mL)

Dextromethorphan HBr 5 mg..... Cough suppressant
Phenylephrine HCl 2.5 mg..... Nasal decongestant

Purposes

- temporarily relieves these symptoms due to the common cold, hay fever, or other upper respiratory allergies:
- cough
- nasal congestion
- sinus congestion and pressure

Uses

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Warnings

- Ask a doctor before use if the child has:
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)
- a sodium-restricted diet

When using this product do not exceed recommended dose

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever
- cough gets worse or lasts for more than 7 days
- cough tends to come back or occurs with fever, rash or headache that lasts

These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)



How can we help?
1-888-217-2117
sudafed.com

NDC 50580-784-04

children's SUDAFED[®] PE


non-drowsy

NASAL DECONGESTANT

Phenylephrine HCl Oral Solution
Nasal Decongestant

RELIEF OF

✔ **Stuffy Nose**
✔ **Sinus Pressure**




Berry
FLAVOR LIQUID

children's SUDAFED[®] PE

non-drowsy

NASAL DECONGESTANT

Drug Facts	
Active ingredient (in each 5 mL)	Purpose
Phenylephrine HCl 2.5 mg.....	Nasal decongestant
Uses	
<ul style="list-style-type: none"> ■ temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies ■ temporarily relieves sinus congestion and pressure 	
Warnings	
<p>Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.</p>	
Ask a doctor before use if the child has	
<ul style="list-style-type: none"> ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ a sodium-restricted diet 	
When using this product do not exceed recommended dose	
Stop use and ask a doctor if	
<ul style="list-style-type: none"> ■ nervousness, dizziness, or sleeplessness occur 	



Other information

- each 5 mL contains: sodium 14 mg
- store between 20-25°C (68-77°F). Protect from light. Store in outer carton until contents are used.
- do not use if carton tape or bottle wrap imprinted with "Sealed For Your Safety" is broken or missing

Inactive ingredients


anhydrous citric acid, carboxymethylcellulose sodium, edetate disodium, FD&C red no. 40, flavors, glycerin, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments?

call 1-888-217-2117 (toll-free) or 215-273-8755 (collect)



3 0045-0537-04 1



available behind
the counter

**Other Products Formulated To
Relieve Your Child's Symptoms**

Tips for Relieving Stuffy Nose Symptoms

Make sure they drink lots
of fluids, especially warm
drinks

Show them how to breathe in
moist air from a hot shower,
humidifier or vaporizer

When treating your child's
symptoms, be sure to use
the dosing cup to properly
dose medication



SNAP THIS
CODE OR
TEXT TYKID
TO 87715

MSG & DATA RATES
MAY APPLY
TEXT HELP FOR HELP
TEXT STOP FOR STOP



How can we help?
1-800-458-1635

NDC 50580-299-01

Children's **TYLENOL** **COLD + FLU**

Acetaminophen, Pain Reliever-Fever Reducer
Chlorpheniramine Maleate, Antihistamine
Dextromethorphan HBr, Cough Suppressant
Phenylephrine HCl, Nasal Decongestant

Oral Suspension

Ages 6-11 Years

- ✓ FEVER
- ✓ SORE THROAT
- ✓ SNEEZING
- ✓ RUNNY NOSE
- ✓ COUGH
- ✓ STUFFY NOSE



Children's **TYLENOL** **COLD + FLU**

Ages 6-11 Years

• Ibuprofen Free • Alcohol Free • Aspirin Free

DIRECTIONS:

STEP 1.

Shake well
before using



STEP 2.

Remove child
protective
cap




STEP 3.

Squeeze your
child's dose
into the
dosing cup



IMPORTANT:



Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: acetaminophen may cause severe skin reactions.

Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- to make a child sleepy
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- if your child is allergic to any of the ingredients in this product

Ask a doctor before use if your child has

- liver disease ■ heart disease ■ high blood pressure
- thyroid disease ■ diabetes
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as chronic bronchitis
- glaucoma

Ask a doctor or pharmacist before use if your child is

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not exceed recommended dose (see overdose warning)
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur

Directions

- this product does not contain directions or complete warnings for adult use.
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 36	under 4 years	do not use
36-47	4 to 5 years	do not use unless directed by a doctor
48-95	6 to 11 years	10 mL

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- store between 20-25°C (68-77°F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "TYLENOL" is broken or missing

Inactive ingredients

anhydrous citric acid, D&C red no. 33, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, microcrystalline cellulose and carboxymethylcellulose sodium, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

Questions or comments?

call 1-800-458-1635 (toll-free) or 215-273-8755 (collect)

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**

McNeil Consumer Healthcare Division

30037519

Fort Washington, PA 19034 USA

©J&JCI 2016

www.tylenol.com



Care To Recycle™



SNAP THIS
CODE OR
TEXT TYKID
TO 87715

MSG & DATA RATES
MAY APPLY
TEXT HELP FOR HELP.
TEXT STOP FOR STOP.



How can we help?
1-800-458-1635

NDC 50580-456-01

Children's **TYLENOL** **COLD + FLU**

Acetaminophen, Pain Reliever-Fever Reducer
Chlorpheniramine Maleate, Antihistamine
Dextromethorphan HBr, Cough Suppressant
Phenylephrine HCl, Nasal Decongestant

Oral Suspension

Ages 6-11 Years

- ✓ FEVER
- ✓ SORE THROAT
- ✓ SNEEZING
- ✓ RUNNY NOSE
- ✓ COUGH
- ✓ STUFFY NOSE



Children's **TYLENOL** **COLD + FLU**

Ages 6-11 Years

• Ibuprofen Free • Alcohol Free • Aspirin Free

DIRECTIONS:

STEP 1.

Shake well
before using



STEP 2.

Remove child
protective
cap





STEP 3.

Squeeze your
child's dose
into the
dosing cup



IMPORTANT:



Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: acetaminophen may cause severe skin reactions.

Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- to make a child sleep
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- if your child is allergic to any of the ingredients in this product

Ask a doctor before use if your child has

- liver disease ■ heart disease ■ high blood pressure
- thyroid disease ■ diabetes
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as chronic bronchitis
- glaucoma

Ask a doctor or pharmacist before use if your child is

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not exceed recommended dose (see overdose warning)
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur

any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use.
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 36	under 4 years	do not use
36-47	4 to 5 years	do not use unless directed by a doctor
48-95	6 to 11 years	10 mL

* or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- store between 20-25°C (68-77°F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "TYLENOL" is broken or missing

Inactive ingredients

anhydrous citric acid, D&C red no. 33, FD&C red no. 40, flavors, glycerin, microcrystalline cellulose and carboxymethylcellulose sodium, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

Questions or comments?

call 1-800-458-1635 (toll-free) or 215-273-8755 (collect)

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**

McNeil Consumer Healthcare Division

30037517

Fort Washington, PA 19034 USA

©J&JCI 2016 www.tylenol.com



Care To Recycle™